

医薬品を正しく購入するための

# 説 明 文 書

第2類医薬品

使用前には必ず添付文書を読んで下さい

| 1         | 名称                    | 太田漢方胃腸薬Ⅱ   |        |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
|-----------|-----------------------|--|--------|--|--|----|-----|--------|-----------|---------|----------|------------|--------|-----------|---------|-----------|--|-------|--------|-------|--------|-----|--------|-----|--------|-------|--------|-------|--------|-----|--------|-----|--------|-------|--------|-------|--------|-------|--------|-------|--------|------|--------|------|--------|--------|--------|--------|--------|
| 2         | 成分・分量                 | <div>3包（1日量：3.0g）中</div> <table><tr><th>成分</th><th>分量</th><th>成分</th><th>分量</th></tr><tr><td>安中散加茯苓末</td><td>2,100 mg</td><td>安中散料加茯苓エキス</td><td>270 mg</td></tr><tr><td colspan="2">（下記生薬の混合末</td><td colspan="2">（下記生薬のエキス</td></tr><tr><td>ブクリョウ</td><td>500 mg</td><td>ブクリョウ</td><td>810 mg</td></tr><tr><td>ケイヒ</td><td>300 mg</td><td>ケイヒ</td><td>486 mg</td></tr><tr><td>エンゴサク</td><td>300 mg</td><td>エンゴサク</td><td>486 mg</td></tr><tr><td>ボレイ</td><td>300 mg</td><td>ボレイ</td><td>486 mg</td></tr><tr><td>ウイキョウ</td><td>200 mg</td><td>ウイキョウ</td><td>324 mg</td></tr><tr><td>シュクシャ</td><td>200 mg</td><td>シュクシャ</td><td>324 mg</td></tr><tr><td>カンゾウ</td><td>200 mg</td><td>カンゾウ</td><td>324 mg</td></tr><tr><td>リョウキョウ</td><td>100 mg</td><td>リョウキョウ</td><td>162 mg</td></tr></table> |        |  |  | 成分 | 分量  | 成分     | 分量        | 安中散加茯苓末 | 2,100 mg | 安中散料加茯苓エキス | 270 mg | （下記生薬の混合末 |         | （下記生薬のエキス |  | ブクリョウ | 500 mg | ブクリョウ | 810 mg | ケイヒ | 300 mg | ケイヒ | 486 mg | エンゴサク | 300 mg | エンゴサク | 486 mg | ボレイ | 300 mg | ボレイ | 486 mg | ウイキョウ | 200 mg | ウイキョウ | 324 mg | シュクシャ | 200 mg | シュクシャ | 324 mg | カンゾウ | 200 mg | カンゾウ | 324 mg | リョウキョウ | 100 mg | リョウキョウ | 162 mg |
| 成分        | 分量                    | 成分   | 分量     |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| 安中散加茯苓末   | 2,100 mg              | 安中散料加茯苓エキス   | 270 mg |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| （下記生薬の混合末 |                       | （下記生薬のエキス  |        |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| ブクリョウ     | 500 mg                | ブクリョウ  | 810 mg |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| ケイヒ       | 300 mg                | ケイヒ  | 486 mg |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| エンゴサク     | 300 mg                | エンゴサク  | 486 mg |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| ボレイ       | 300 mg                | ボレイ  | 486 mg |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| ウイキョウ     | 200 mg                | ウイキョウ  | 324 mg |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| シュクシャ     | 200 mg                | シュクシャ  | 324 mg |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| カンゾウ      | 200 mg                | カンゾウ   | 324 mg |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| リョウキョウ    | 100 mg                | リョウキョウ   | 162 mg |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| 3         | 用法・用量                 | <div>次の量を食間（就寝前を含む）又は空腹時に服用してください。</div> <table><tr><th>年齢</th><th>1回量</th><th>1日服用回数</th></tr><tr><td>成人(15歳以上)</td><td>1包</td><td rowspan="2">3回</td></tr><tr><td>7～14歳</td><td>2/3包</td></tr><tr><td>7歳未満</td><td colspan="2">服用しないこと</td></tr></table> <div>（食間とは食後2～3時間のことをいいます。）</div>   |        |  |  | 年齢 | 1回量 | 1日服用回数 | 成人(15歳以上) | 1包      | 3回       | 7～14歳      | 2/3包   | 7歳未満      | 服用しないこと |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| 年齢        | 1回量                   | 1日服用回数   |        |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| 成人(15歳以上) | 1包                    | 3回   |        |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| 7～14歳     | 2/3包                  |  |        |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| 7歳未満      | 服用しないこと               |  |        |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| 4         | 効能・効果                 | 体力中等度以下で、腹部は力がなくて、神経過敏で胃痛又は腹痛があつて、ときに胸やけや、げっぷ、胃もたれ、食欲不振、はきけ、嘔吐などを伴うものの次の諸症： 神経性胃炎、慢性胃炎、胃腸虚弱  |        |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| 5         | 保健衛生上の危害を防止するために必要な事項 | <div>次に該当する人はお申し出ください</div> <ul style="list-style-type: none"><li>・ 医師の治療を受けている人。</li><li>・ 妊婦又は妊娠していると思われる人。</li><li>・ 今までに薬により発疹・発赤、かゆみ等を起こしたことがある人。</li></ul>   |        |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |
| 6         | 薬剤師または登録販売者が必要と判断する事項 |  |        |  |  |    |     |        |           |         |          |            |        |           |         |           |  |       |        |       |        |     |        |     |        |       |        |       |        |     |        |     |        |       |        |       |        |       |        |       |        |      |        |      |        |        |        |        |        |

## 〔注意事項〕

1. 本紙の内容は、お客様が医薬品を購入・選択時に、役立たせるために必要な情報です。
2. 情報提供が不必要とされるお客様には、申し出により情報提供は行いません。
3. 使用後、体調に変化等があった場合（副作用など）には使用を中止し、すぐに購入された店舗の薬剤師または登録販売者にご相談下さい。

## OHTA KANPO ICHOYAKU II

### 【Features】

#### 1. *Efficacious on nervous gastritis and gastric atonia*

OHTA KANPO ICHOYAKU II takes excellent effect on nervous gastritis and gastric atonia which usually attack those who suffer from serious stress through the pressure of work or those who are nervous and get tired easily.



Poria Sclerotium has tranquilizing effect and is quite efficacious on nervous gastritis, interacting with other ingredients such as Cinnamon bark, Amomum seed, Alpinia officinarum Rhizome, Fennel and Oyster shell.

#### 2. *Powerful effect on chronic gastritis*

OHTA KANPO ICHOYAKU II exerts excellent effect on the following symptoms which are caused by chronic gastritis: usual feeling of stomach weakness, lost appetite and frequent gastrointestinal pain.

On chronic gastritis with stomach pain work Corydalis Tuber, Alpinia officinarum Rhizome, Glycyrrhiza and Oyster shell. Similarly, on chronic gastritis with lost appetite, heartburn, belch and nausea operate Amomum seed, Fennel, Cinnamon bark and Alpinia officinarum Rhizome.

#### 3. *Easily to take*

OHTA KANPO ICHOYAKU II is powder dosage, which dissolves immediately and is easily to take. It is due to control the evaporation of essential oil contained in crude drugs and avoids an unnecessary process as much as possible.

### 【Directions】

Over 15 years old: 1 sachet

From 7 to 14 years old: 2/3 sachet

Take 3 times a day between meals (include at bedtime) or on an empty stomach.

Also available such tablet as “OHTA KANPO ICHOYAKU II <TABLET>”

Over 15 years old: 3 tablets

From 7 to 14 years old: 2 tablets

Take 3 times a day between meals (include at bedtime) or on an empty stomach.

※Children: Not to be given to infants under 7 years.

## 【Indications】

The following symptoms in thin persons who tend to have diminished abdominal muscle tension and who have stomachache or abdominal pain and occasionally heartburn, belching, loss of appetite, and nausea: Nervous gastritis, chronic gastritis and gastric atonia.

## 【Ingredients】

Anchu-San plus Poria Sclerotium (2,100mg)

Poria Sclerotium 500mg Corydalis tuber 300mg Fennel 200mg Glycyrrhiza 200mg  
Cinnamon bark 300mg Oyster shell 300mg Amomum seed 200mg  
Alpinia officinarum Rhizome 100mg  
Mixed powder of the above herbal medicines

Anchu-San plus Poria Sclerotium extract (270mg)

Poria Sclerotium 810mg Corydalis tuber 486mg Fennel 324mg Glycyrrhiza 324mg  
Cinnamon bark 486mg Oyster shell 486mg Amomum seed 324mg  
Alpinia officinarum Rhizome 162mg  
Extract of the above herbal medicines

## 【Consult】

1. The following persons should consult a physician or pharmacist before using the medication;
  - (1) Those who are under treatment by a physician.
  - (2) Those who are pregnant or could be pregnant.
  - (3) Those who have ever been rash, redness and itching caused by medicines.
2. Discontinue the medication immediately when the following cases have occurred and consult a physician or pharmacist, showing this Package Insert;
  - (1) When the following symptoms have occurred after taking the medication:  
Areas concerned: Skin      Symptoms: Rash, redness, itching
  - (2) When symptoms have not improved ever after medication of one month.

Manufacturer; OHTA' S ISAN CO., LTD.  
3-2, Sengoku 2-Chome, Bunkyo-Ku  
Tokyo, Japan